

Synonym

IL7Ra,CD127

Source

Human IL-7 R alpha Protein, His Tag(IL7-H52H7) is expressed from human 293 cells (HEK293). It contains AA Glu 21 - Asp 239 (Accession # [P16871-1](#)).

Molecular Characterization

IL-7 R alpha(Glu 21 - Asp 239)
P16871-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 27.3 kDa. The protein migrates as 38-48 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human IL-7 R alpha Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).


Bioactivity-ELISA

Discounts, Gifts,
and more!



Human IL-7 R alpha / CD127 Protein, His Tag

Catalog # IL7-H52H7



Immobilized Human IL-7 R alpha Protein, His Tag (Cat. No. IL7-H52H7) at 5 µg/mL (100 µL/well) can bind Human IL-7, Fc Tag (Cat. No. IL7-H5253) with a linear range of 0.039-0.625 µg/mL (QC tested).

Background

Interleukin-7 receptor subunit alpha (IL7Ra) is also known as CD antigen CD127, which belongs to the type I cytokine receptor family and type 4 subfamily. IL7Ra /CD127 contains 1 fibronectin type-III domain. IL7Ra /CD127 is expressed on various cell types, including naive and memory T cells and many others. IL7Ra /CD127 is receptor for interleukin-7 and also acts as a receptor for thymic stromal lymphopoietin (TSLP). The IL7 receptor is a heterodimer of IL7R and IL2RG. The TSLP receptor is a heterodimer of CRLF2 and IL7R.

