

Synonym

IL7Ra,CD127

Source

Biotinylated Human IL-7 R alpha Protein, Fc,Avitag(IL7-H82F9) is expressed from human 293 cells (HEK293). It contains AA Glu 21 - Asp 239 (Accession # [P16871-1](#)).

Molecular Characterization

IL-7 R alpha(Glu 21 - Asp 239) P16871-1	Fc(Pro 100 - Lys 330) P01857	Avi
--	---------------------------------	-----

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™). The protein has a calculated MW of 53.5 kDa. the protein migrates as 67-75 kDa under reducing (R) condition, and >116 kDa under non-reducing (NR) condition (SDS-PAGE).

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Human IL-7 R alpha Protein, Fc,Avitag on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA





Immobilized Human IL-7, Fc Tag (Cat. No. IL7-H5253) at 2 µg/mL (100 µL/well) can bind Biotinylated Human IL-7 R alpha Protein, Fc,Avitag (Cat. No. IL7-H82F9) with a linear range of 1-20 ng/mL (QC tested).

Background

Interleukin-7 receptor subunit alpha (IL7Ra) is also known as CD antigen CD127, which belongs to the type I cytokine receptor family and type 4 subfamily. IL7Ra /CD127 contains 1 fibronectin type-III domain. IL7Ra /CD127 is expressed on various cell types, including naive and memory T cells and many others. IL7Ra /CD127 is receptor for interleukin-7 and also acts as a receptor for thymic stromal lymphopoietin (TSLP). The IL7 receptor is a heterodimer of IL7R and IL2RG. The TSLP receptor is a heterodimer of CRLF2 and IL7R.

